

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

PATENT SPECIFICATION

(11) 1 571 683

1 571 683

(21) Application No. 6009/76 (22) Filed 16 Feb. 1976
 (21) Application Nos. 27301/76 and 27302/76
 (22) Filed 30 June 1976
 (23) Complete Specification filed 15 Feb. 1977
 (44) Complete Specification published 16 July 1980
 (51) INT CL¹ C07D 501/34; A61K 31/545
 (52) Index at acceptance
 C2C 1314 1470 214 215 246 247 253 256 25Y 28X 302 30Y
 33Y 340 342 346 34Y 351 352 357 366 368 638 801
 80Y AA KE



(72) Inventors MICHAEL GREGSON and RICHARD BROOK
 SYKES

(54) ESTER DERIVATIVES OF CEFUROXIME

(71) We, GLAXO OPERATIONS U.K. LIMITED, formerly known as Glaxo Laboratories Limited, a British company, of Greenford, Middlesex, do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—
 This invention is concerned with improvements in or relating to cephalosporin antibiotics. More particularly the invention is concerned with biologically acceptable ester derivatives of (6R,7R) - 3 - carbamoyloxy-methyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2-methoxyiminoacetamido]ceph - 3 - em - 4-carboxylic acid (i.e. the *syn* isomer), which has the approved name "cefuroxime".
 Cefuroxime, as disclosed in British Patent Specification No. 1453049 is a valuable broad spectrum antibiotic characterised by high activity against a wide range of gram-positive and gram-negative microorganisms, this property being enhanced by the very high stability of the compound to β -lactamases produced by a range of gram-negative microorganisms. Additionally the compound is stable in the body owing to its resistance to the action of mammalian esterases; and gives high serum levels following parenteral administration (e.g. in the form of the sodium salt) to human and animal subjects, while exhibiting low serum binding.
 Cefuroxime and its salts, for example alkali metal salts such as the sodium salt, are principally of value as injectable antibiotics since they are poorly absorbed from the gastro-intestinal tract and are therefore present in sera and urine only in low concentrations after oral administration. We have accordingly conducted extensive studies into the possible activity upon oral administration of various derivatives of cefuroxime, since the development of derivatives which are absorbed

through the gastro-intestinal tract and exhibit good antibacterial activity following oral administration would extend still further the valuable therapeutic potential of cefuroxime.
 It is known from the literature pertaining to β -lactam antibiotics that the effect upon oral administration of penicillin antibiotics such as ampicillin can be improved by converting the carboxy group at the 3-position of the penam nucleus to certain esterified carboxy groups; there have also been some proposals that the activity upon oral administration of certain cephalosporin antibiotics may be enhanced by esterification in similar manner. It is believed that the presence of an appropriate esterifying group enhances absorption of the compound from the gastro-intestinal tract, whereupon the esterifying group is hydrolyzed by enzymes present in, for example, serum and body tissues to yield the antibiotically active parent acid. It will be appreciated that the precise nature of the esterifying group is critical since it is necessary that the ester should be sufficiently stable to allow the ester to reach the site of absorption without undergoing significant degradation, e.g. in the stomach, while on the other hand the ester must be sufficiently susceptible to esterase hydrolysis so that the antibiotically active parent acid is liberated within a short time of the ester being absorbed.
 The selection of a particular esterifying group to enhance the effect upon oral administration of a β -lactam antibiotic will also be influenced by the specific β -lactam compound chosen. Thus, for example, esterifying groups which have been found effective in improving the activity of orally administered penicillin antibiotics do not necessarily convey similar advantages to antibiotics of the cephalosporin series. An example which may be cited here is the case of pivaloyloxymethyl esters. Thus, the pivaloyloxymethyl ester of, for example,

ERRATA

SPECIFICATION No. 1,571,683


- Page 1, line 56, *after* may insert be
Page 2, line 62, for amoyloxymethyl read amoyloxymethyl
Page 2, line 67, for methoxyiminoacetamide read methoxyiminoacetamido
Page 2, line 74, for omyloxymethyl read amoyloxymethyl
Page 2, line 78, for (fur- -yl) read (fur-2-yl)
Page 2, line 88, *after* shown insert by
Page 3, line 26, for of read or
Page 3, line 46, *after* a insert 1,
Page 3, line 56, *before* as insert such
Page 3, line 62, for effectde read effected
Page 5, line 19, *after* yl)- insert 2-
Page 5, line 33, *after* (EtOH) delete —
Page 5, line 45, for N,N-dimethylformadie read N,N-dimethylformamide
Page 5, line 46, for bromobutyrate read bromoethylbutyrate
Page 5, line 91, *before* fur insert (
Page 5, line 107, for carbaomyloxymethyl read carbamoyloxymethyl
Page 6, line 13, *before* 7 delete (
Page 6, line 17, for -1- read -7-
Page 6, line 41, *after* nm insert (E
Page 6, line 44, *after* 5.95% insert]
Page 6, line 64, *after* nm insert (E
Page 6, line 71, *after* (R and S)- insert 1-
Page 6, line 107, *before* Acetoxyethyl delete (
Page 6, line 113, *after* (fur-2-yl) delete -2)- insert -2-
Page 7, line 14, for -2- read -3-
Page 7, line 60, *after* (EtOH insert)
Page 7, line 70, *after* 5.9 delete — insert %
Page 7, line 71, for Isober read Isomer
Page 12, TABLE 1 (Continued) heading to Column 3, for HN₂ + NH read NH₂ + NH
Page 13, line 36, for -(2- read -2- (
Page 13, line 57, for sealed read sealed
Page 13, line 96, *after* (Z) insert - *after* (fur-2-yl)- insert 2-
Page 14, line 12, for of read or
Page 14, line 14, for methoxyiminoacetate read methoxyiminoacetic for reaction read reactive

THE PATENT OFFICE

15th December, 1980

ampicillin is known to improve the oral absorption of ampicillin. The pivaloyloxymethyl ester of cefuroxime, on the other hand, exhibits little effect upon oral administration, possibly because the ester is not sufficiently absorbed from the gastro-intestinal tract or alternatively is substantially resistant to esterase hydrolysis so that the antibiotically active acid is not liberated to any significant extent following absorption.

We have now found that esters of cefuroxime, which may be represented by the formula



(wherein R¹ is a primary or secondary alkyl group containing 1 to 4 carbon atoms and R² is a primary or secondary alkyl group containing 1 to 6 carbon atoms provided that at least one of the groups R¹ and R² is methyl, and the asterisk denotes an asymmetric carbon atom) possess properties which render these compounds of significant potential value as orally administrable antibiotics. The individual diastereoisomers, as well as mixtures thereof, are embraced by the invention.

The references herein to primary or secondary alkyl groups denote alkyl groups in which the carbon atom of attachment is bonded to at least one hydrogen atom.

Examples of compounds of formula I include those wherein R¹ is a methyl group and R² is an alkyl group containing 2 to 4 carbon atoms and those wherein R² is a methyl group and R¹ is a primary or secondary alkyl group containing 1 to 4 carbon atoms.

The esters (I) possess reasonable stability as evidenced by the fact that they exhibit low antibacterial activity *in vitro* compared to cefuroxime (this indicates that a high proportion of ester remains unchanged throughout the *in vitro* tests and so confirms the stability of the esters). The esters are, on the other hand, extremely susceptible to esterase hydrolysis leading to formation of cefuroxime, as evidenced by *in vitro* tests employing esterases derived from rat liver, human liver and human serum.

In vivo testing in rats confirms that oral administration of the esters (I) leads to significantly greater absorption of cefuroxime, as evidenced by higher serum levels and increased urinary recovery, than does oral administration of cefuroxime itself.

Of the compounds of formula (I) the following have been shown to provide particularly good absorption of cefuroxime:—

1 - acetoxyethyl (6R,7R) - 3 - carbamoyl-

oxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate;

1 - propionyloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate;

1 - butyryloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamide]ceph - 3 - em - 4 - carboxylate;

1 - isovaleryloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate;


1 - acetoxyheptyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate;

1 - acetoxybutyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate;

1 - acetoxypropyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.


The first compound mentioned above is particularly preferred, since it provides especially high absorption of cefuroxime as shown *in vivo* testing in mice, rats and dogs.

The compounds (I) may be prepared in conventional manner, for example by reacting cefuroxime or a salt thereof (e.g. an alkali metal salt such as the sodium or potassium salt or an onium salt, e.g. an ammonium salt for example a quaternary ammonium salt) with a haloester of formula



(where R¹, R² and the asterisk have the above-defined meanings and X is halogen such as chlorine, bromine or iodine). The reaction is conveniently effected in solution in an inert organic solvent (e.g. an N,N-disubstituted amide such as N,N-dimethylformamide or N,N-dimethylacetamide, a ketone such as acetone, a sulfoxide such as dimethylsulphoxide, a nitrile such as acetonitrile, or hexamethylphosphoric triamide) at a temperature in the range -50° to +150°C, e.g. -10° to +50°C, conveniently between 0°C and room temperature. When a cefuroxime salt, for example, the potassium salt, is employed as starting material and the reaction is effected in a nitrile solvent, a crown ether such as 18-crown-6 may, if desired, be employed. When cefuroxime is employed as starting material it may be advantageous to effect the reaction in the presence of a base, e.g. a weak inorganic

The esters (I) may also be prepared by acylation of a compound of formula:—



50 The free acid may itself be used as the acylating agent. Such acylations are desirably conducted in the presence of, for example, a carbodiimide such as N,N' -dicyclohexylcarbodiimide; a carbonyl compound such as
55 carbonyldiimidazole; or an oxazolanium salt as N -ethyl-5-phenylisoxazolanium-3'-

The compositions may contain from 0.1% upwards, e.g. 0.1—99%, conveniently from 10—60% of the active ingredient (I), depending on the method of administration. Compositions in dosage unit form conveniently contain 50—500 mg of the active ingredient (calculated as cefuroxime). Doses employed for adult human treatment will typically be in the range 500—5000 mg per day, e.g. 1500 mg per day, (calculated as cefuroxime).

although the precise dose will depend on, *inter alia*, the frequency of administration.

The following Examples illustrate the invention. All temperatures are in °C. The melting points were determined by the capillary method and are uncorrected. Those prefixed (M_y^x) where x is the rate of heating (in °C per minute) and y is the insertion temperature were measured in a Mettler (registered Trade Mark) apparatus. In Example 5 the potassium carbonate employed was dried at 120° *in vacuo* and finely ground. The *N,N*-dimethylformamide employed was dried and purified by passage through acidic alumina.

HPLC stands for "high pressure liquid chromatography". Detection was achieved by ultraviolet light at 276 nm. Relative peak areas were measured at this wave length. (The λ_{max} of the desired compound occurs at 276 to 277 nm).

The n.m.r. spectra figures for the products of the Examples 1 to 7 given in Table 1 hereinafter indicate that the compounds are obtained as approximately 1:1 mixtures of the R and S isomers.

Preparation 1.

(R,S) 1-Bromoethyl propionate.

Acetaldehyde (1.7 ml, 1.34 g, 30.5 mmole) was added dropwise with stirring to propionyl bromide (3.18 g, 23.2 mmole) (0–5°C). The mixture was allowed to warm up to room temperature (ca 20°) over 1 hour. The product was distilled to give the title ester (2.7 g) as a liquid b.p. 41 to 50°/15 mm which was characterised by its nmr ($CDCl_3$) and infrared ($CHBr_3$) spectra.

Preparation 2.

(R,S) 1-Bromoethyl n-butyrate.

Acetaldehyde (2 ml) was added dropwise with stirring to n-butyryl bromide (2.09 g, 13.8 mmole) at 0–10°C. After the initial reaction the mixture was stored at ca 4° for 2 days. The product was distilled to give the title ester (1.87 g) as a liquid b.p. 63 to 65°/14 mm which was characterised by its nmr ($CDCl_3$) and infrared ($CHBr_3$) spectra.

Preparation 3.

(R,S) 1-Bromoethyl 3-methylbutyrate.

Acetaldehyde (2.3 ml, 1.8 g, 41 mmole) was added dropwise with stirring to 3-methylbutyryl bromide (5.33 g, 32 mmole) at 0°C. The mixture was allowed to warm up to room temperature (ca. 20°C) over half an hour.

The title ester was isolated in two fractions (2.99 g and 1.64 g) by distillation; b.p.'s 60 to 70°/25 mm and 70 to 75°/25 mm respectively.

Both fractions were characterised by their nmr ($CDCl_3$) spectra. The second fraction was the purer by nmr.

Preparation 4.

(R,S) 1-Bromo-n-butyl acetate.

To cooled (ca 0°) acetyl bromide (1.49 ml, 20 mmole) was added n-butyraldehyde (1.76 ml, 20 mmole). The reaction mixture was allowed to warm up to room temperature over 1 hour to give a pale brown solution. This was distilled *in vacuo* to give two fractions:—

(i) bp 60 to 70°/27 mm (0.78 g) 70

(ii) bp 70 to 80°/27 mm (1.64 g)

Fraction (ii) contained the title ester which was characterised by its nmr ($CDCl_3$) and infrared ($CHBr_3$) spectra.

Preparation 5.

(R,S) 1-Bromopropyl acetate.

The preparation was analogous to Preparation 4 except that the following reagents were substituted and the mixture was allowed to react overnight at ca 5°.

Acetyl bromide (1.5 ml, 20 mmole)

Propionaldehyde (2.2 ml, 31 mmole).

The title ester (1.56 g) was isolated as a liquid bp 50 to 60°/20 mm characterised by its nmr ($CDCl_3$) spectrum.

Example 1.

(R and S) - 1 Acetoxyethyl (6R,7R) - 3-carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

Potassium carbonate (760 mg, 5.5 mmole) was added to a solution of (6R,7R) - 3-carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylic acid (4.57 g, 11 mmole) in *N,N*-dimethylformamide (25 ml) and the mixture was stirred at ca 20° for 25 minutes. 1-Bromoethyl acetate (1.8 g, 11 mmole) in *N,N*-dimethylformamide (5 ml) was added to the above solution and the reaction mixture was stirred for 40 minutes at ca 20°. The reaction mixture was worked up by pouring it into excess 2N hydrochloric acid, followed by extraction with ethyl acetate (3 times). The combined organic extracts were washed with 2N hydrochloric acid and saturated sodium bicarbonate solution, dried (magnesium sulphate) and evaporated *in vacuo* to yield a foam which was dissolved in ethyl acetate and precipitated from diethyl ether.

The resulting precipitate was filtered off and dried to give the title compound (780 mg).

The mother liquors were evaporated to a foam which was dissolved in ethyl acetate and precipitated from di-isopropyl ether to give a further crop of the title compound (1.21 g). This sample was dried *in vacuo* for 2 days at 22° in order to remove di-isopropyl ether. The physical constants of the second crop of the title compound are:—mp (M_{10}^2)

72°; $[\alpha]_D + 84^\circ$ (c 0.87, DMSO); λ_{\max} 18,850; [Found:—C, 48.8; H, 5.0; N, 10.1; S, 5.5; $C_{22}H_{28}N_4O_{10}S$ (538.5) requires C, 49.1; H, 4.9; N, 10.4; S, 5.95%]. The infrared and nmr data are shown in Table 1 hereinafter.

Example 2.

(R and S) - 1 - Propionyloxyethyl (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

The method of preparation was analogous to that described in Example 1; 1-bromoethyl propionate (1.5 g, 8.3 mmole) was reacted with potassium (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (3.8 g, 8.3 mmole) in N,N-dimethylformamide (25 ml) at ca 22° for 30 minutes.

The reaction mixture was worked up as described in Example 1 to yield an oil which was dissolved in ethyl acetate and precipitated from diethyl ether (250 ml). The filtrate was evaporated to an oil which was dissolved in ethyl acetate and added dropwise to di-isopropyl ether. The precipitate was filtered and washed with di-isopropyl ether and dried to give the title compound (1.46 g, 2.8 mmole), mp (M_{150}°) 81°; $[\alpha]_D + 66^\circ$ (c 1.2, DMSO); λ_{\max} (EtOH)-276.5 nm (E 1% 362, ϵ 18,985). The infrared and nmr data are shown in Table 1 hereinafter.

Example 3.

(R and S) - 1 - Butyryloxyethyl (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

A solution of potassium (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (1.025 g, 2.22 mmole) in N,N-dimethylformamide (12 ml) was treated with 1-bromobutyrate (649 mg, 3.33 mmole) in N,N-dimethylformamide (2 ml) for 30 minutes at room temperature.

The reaction mixture was then worked up as described in Example 1 and after drying (over magnesium sulphate) and evaporation, a pale yellow foam (964 mg) was obtained. The foam was triturated with cyclohexane (80 ml) and the resultant solid was filtered off and washed with cyclohexane (2 x 20 ml) and dried *in vacuo* to give the title compound (711 mg) as an off-white solid; mp (M_{150}°) 84°; $[\alpha]_D + 36^\circ$ (c 1.13, DMSO); λ_{\max} (EtOH)-276.5 nm (E 1% 350, ϵ 1 cm

18,850); [Found:—C, 48.8; H, 5.0; N, 10.1; S, 5.5; $C_{22}H_{28}N_4O_{10}S$ (538.5) requires C, 49.1; H, 4.9; N, 10.4; S, 5.95%].

The infrared and nmr data are shown in Table 1 hereinafter.

Example 4(a).

(R and S) - 1 - Isovaleryloxyethyl (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate and (R and S) - 1 - Isovaleryloxyethyl (4R, 6R, 7R) - 3 - carbamoyloxymethyl - 1 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 2 - em - 4 carboxylate.

1 - Bromoethyl - 3 - methylbutyrate (2.38 g, 11.4 mmole) was added to a stirred solution of potassium (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (4.2 g, 9.1 mmole) in N,N-dimethylformamide (50 ml) at ca 22° for 35 minutes. The reaction mixture was then worked up in a similar manner to that described in Example 1 and the product (2.15 g) was precipitated from di-isopropyl ether. The nmr spectrum (DMSO-d₆) indicated that the product was approximately a 1:1 mixture of the two title compounds.

(b) (R and S) - 1 - Isovaleryloxyethyl (1S, 6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate 1 - oxide.

A solution of the mixture of esters produced in Example 4(a) (2.0 g) in dichloromethane (50 ml) was treated with m-chloroperbenzoic acid (1.0 g, 5.7 mmole) for 40 minutes at ca 22°.

The solvent was evaporated *in vacuo* and the residue was triturated with diethyl ether to give the title compound (1.8 g) as a solid, mp (M_{150}°) 167.5°; $[\alpha]_D + 52^\circ$ (c 0.5, DMSO); [Found; C, 47.1; H, 4.7; N, 10.05; S, 5.8; $C_{21}H_{27}N_4O_{11}S$ (568.5) requires C, 48.6; H, 4.95; N, 9.85; S, 5.65%].

(c) (R and S) - 1 - Isovaleryloxyethyl (6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

Potassium iodide (1.95 g, 11.7 mmole) and acetyl chloride (460 mg, 5.9 mmole) were added successively to a solution of (R and S) - 1 - isovaleryloxyethyl (1S, 6R, 7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate 1-oxide (1.67 g, 2.9 mmole) in N,N-dimethylformamide (50 ml).

The solution was stirred at ca 22° for 35 minutes and then added dropwise to an aqueous solution of sodium metabisulphite. The precipitate was filtered off and washed

with water and dried *in vacuo* over phosphorus pentoxide to give the title compound (1.19 g, 2.1 mmole) as a solid, mp (M_{11}^{21}) 92°; $[\alpha]_D + 24^\circ$ (c 0.9, DMSO); λ_{max} (EtOH) 276 nm (E 1% 1 cm 360, ϵ 19,890); [Found; C, 48.1; H, 4.95; N, 10.35; S, 6.0; $C_{22}H_{21}N_4O_{10}S$ (552.5) requires C, 49.9; H, 5.1; N, 10.1; S, 5.8%].

The infrared and nmr data are shown in Table 1 hereinafter.

Example 5.

(R and S) - 1 - Acetoxybutyl (6R,7R) - 3-carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate. Potassium (6R,7R) - 3 - carbamoyloxymethyl - 1 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (1.850 g, 4 mmole) was added to a solution of 1-bromobutyl acetate (780 mg, 4 mmole) in purified N,N-dimethylformamide (10 ml) resulting in the formation of a brown solution and evolution of heat. After ca 10 minutes solid started separating out and after 20 minutes the reaction mixture was worked up by pouring it into 2N hydrochloric acid (120 ml) to give a pale yellow solid which dissolved on addition of ethyl acetate (120 ml).

The organic layer was separated and washed with saturated aqueous sodium bicarbonate (120 ml) and brine (60 ml), dried over magnesium sulphate and evaporated to a pale yellow foam (1.378 g). Trituration of this foam with di-isopropyl ether (30 ml) gave a pale solid which was filtered off and washed with more di-isopropyl ether and dried *in vacuo* to give the title compound (1.261 g) as a cream powder, mp 59 to 68°; $[\alpha]_D^{21} + 54.5^\circ$ (c 1.0, DMSO); λ_{max} (EtOH) 277 nm (E 1% 1 cm 333, ϵ 17,930); [Found; C, 50.1; H, 5.5; N, 9.4; S, 5.1; $C_{22}H_{21}N_4O_{10}S$ (538.5) requires C, 49.05; H, 4.85; N, 10.4; S, 5.95%. The nmr and infrared data are shown in Table 1 hereinafter.

Example 6.

(R and S) - 1 - Acetoxypentyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate. A solution of potassium (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (3.7 g, 8 mmole) in N,N-dimethylformamide (50 ml) was stirred at ca 22° for 45 minutes with 1-bromopentyl acetate (1.45 g, 8 mmole). The workup was similar to that described in Example 5 except that the crude product was purified by precipitation from ethyl

acetate solution using di-isopropyl ether to give on drying *in vacuo* the title compound (920 mg), mp (M_{11}^{21}) 81°; $[\alpha]_D + 69^\circ$ (c 0.87 DMSO); λ_{max} (EtOH) 277 nm (E 1% 1 cm 349, ϵ 18,305); [Found; C, 48.15; H, 4.8; N, 10.45; S, 5.9; $C_{22}H_{21}N_4O_{10}S$ (524.5) requires C, 48.1; H, 4.6; N, 10.7; S, 6.1%].

The nmr and infrared data are shown in Table 1 hereinafter.

Example 7.

(R and S) - Acetoxyheptyl (6R,7R) - 3-carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

Potassium carbonate (0.21 g) was added to a solution of (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylic acid (1.27 g) in N,N - dimethylformamide (7 ml), with stirring at 23°. Most of the potassium carbonate dissolved within 10 minutes, giving a dark solution. 1-Bromoheptyl acetate (0.72 g) was added as a solution in N,N - dimethylformamide (1.5 ml). Precipitation commenced after 15 minutes, and after 18 minutes the reaction mixture was poured into 2N-hydrochloric acid (75 ml), giving a brown gum. This dissolved on addition of ethyl acetate (75 ml). The organic layer was separated, washed successively with 2N-hydrochloric acid (75 ml) and saturated sodium bicarbonate solution (75 ml), and was dried (Mg SO₄) and evaporated *in vacuo* to give a brown glass (1.01 g). Trituration of this material with petroleum ether (b.p. 40 to 60° - 3 x 15 ml) gave a pale yellow solid which was filtered, washed with petroleum ether (40-60°) and dried *in vacuo* to give the title compound as a pale yellow powder (0.75 g), m.p. 68 to 71° (decomp.); $[\alpha]_D^{21} + 46^\circ$ (c 1.00, DMSO); λ_{max} (EtOH) 276.5 nm (c 18,154).

The nmr and infrared data are shown in Table 1 hereinafter.

Example 8.

1ξ - (Acetoxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (Isomer A).

A solution of (R and S) - 1 - Acetoxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - (methoxyiminoacetamido]ceph - 3 - em - carboxylate (ca. 1:1; ca. 1 g) in methanol (3 ml) was cooled to 0° and left overnight to give a crystalline deposit of Isomer A (300 mg) which was shown by nmr spectroscopy (DMSO-d₆) to contain essentially one isomer.

The mother liquors were evaporated to dryness *in vacuo* and the residue was dissolved

in ethyl acetate and precipitated from petroleum (40—60°). Nmr (DMSO- d_6) indicated that the precipitate consisted of a ca. 65:35 mixture of diastereoisomers B and A respectively.

Example 9.

Separation of the diastereoisomers of 1-acetoxyethyl - (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (Isomer A and Isomer B).

A solution of (R and S) - 1 - Acetoxyethyl - (6R,7R) - 2 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate (ca. 1:1; 5.0 g, 9.8 mmole) in methanol (7.5 ml) at ca. 25° was seeded with a sample of Isomer A (prepared as described in Example 8). The solution solidified and was refrigerated at 0° overnight. Filtration gave a solid (1.7 g) which on recrystallisation from methanol (30 ml) afforded essentially pure Isomer A (1.28 g) (Fraction 1).

A second crop of crystals (110 mg, Fraction 2) was obtained from the mother-liquors. This was shown by nmr (DMSO- d_6) to contain a ca. 1:1-ratio of Isomer A to Isomer B.

The residual mother-liquors were evaporated *in vacuo* to dryness and the residual gum was taken up in ethyl acetate and refrigerated whereupon the solution solidified, so ethyl acetate was added to give a total volume of ca. 15 ml and the mixture was heated to reflux. A small portion of solid did not dissolve and so was filtered off (Fraction 3, 190 mg). Fraction 3 was shown by nmr spectro-

scopy (DMSO- d_6) to consist of essentially Isomer B.

The filtrate was refrigerated and slowly solidified. The resultant solid was filtered off and dried (Fraction 4, 410 mg). Fraction 4 consisted of essentially Isomer B (ca 80% pure) as shown by nmr spectroscopy (DMSO- d_6) and HPLC.

The mother liquor from the Fraction 4 separation was evaporated to dryness and the resultant solid triturated with ethyl acetate-diethyl ether. The solid obtained was filtered and dried (Fraction 5, 640 mg). Fraction 5 consisted of an approximately 70:30 mixture of Isomers B to A, as shown by nmr (DMSO- d_6).

Fraction 1 (Isomer A) and Fraction 4 (Isomer B) have the following physical properties:


Isomer A (Fraction 1): m.p. $[M]_{175}^{25}$ 191°, $[\alpha]_D^{25} + 53^\circ$ (c 0.9, DMSO), λ_{max} (EtOH) 277 nm ($E_{1\%}^{1\text{cm}}$ 414, ϵ 21,130), [Found:— C, 46.95; H, 4.4; N, 10.9; S, 6.5, $C_{20}H_{22}N_4O_{10}S$ (510.5) requires C, 47.1; H, 4.3; N, 10.9; S, 6.3%]. HPLC indicated an isomer purity of ca 94%.

Isomer B (Fraction 4): m.p. $[M]_{115}^{25}$ 129°, $[\alpha]_D^{25} + 11^\circ$ (c 1.2, DMSO), λ_{max} (EtOH) 277 nm ($E_{1\%}^{1\text{cm}}$ 422, ϵ 22,700),

[Found: C, 46.8; H, 4.5; N, 10.35; S, 5.9 $C_{20}H_{22}N_4O_{10}S$, 0.3 mole EtOAc (539.9) requires C, 47.45; H, 4.6; N, 10.4; S, 5.9—]. HPLC indicated an Isomer B purity of ca 80%.

The nmr and infrared data for the two isomers are shown in Table 1 hereinafter.

Physical Properties of the Products of Examples 1 to 7 and 9



Example No.	Solvent	τ (100MHz; J Hz)								R	Assignments for R
		a_1, a_2, a_3	b	c	d	e	f	g	h		
1	DMSO-d ₆	2.11(m) 3.2 to 3.4(m)	6.08	0.20 (d9)	4.06 (multi-plets)	4.69 4.73 (d5)	6.23 6.45 (J18)	5.13 5.35 (J12)	3.30 obscured by a_2 and a_1	i O k -CHOCCCH ₃ CH ₃ j	i 2.96, 3.06(m) j 8.49(d6) k 7.92
2	DMSO-d ₆	2.14(m) 3.2 to 3.4(m)	6.09	0.22 (d8)	4.10 (multi-plets)	4.74 (d5)	6.24 6.47 (J18)	5.14 5.38 (J12)	3.36	i O -CHOCCCH ₃ CH ₃ CH ₃ j	i 2.95, 3.04(m) j 8.49(d6) k 7.60(q7) l 8.94(r7)
3	DMSO-d ₆	2.16(m) 3.2 to 3.4(m)	6.11	0.22 (d8)	4.12 4.15 (dd 8,5)	4.75 4.77 (d5)	6.40 (s)	5.16 5.39 (J12)	3.38	i O -CHOCCCH ₃ CH ₃ CH ₃ CH ₃ j	i 2.99, 3.06(q6) j 8.53(d6) k 7.68(r7) l 8.3 to 8.6(m) m 9.12(r7)

TABLE 1 (Continued)

Example No.	Solvent	τ (100MHz; J Hz)								R	Assignments for R
		a_1, a_2, a_3	b	c	d	e	f	g	h		
4	DMSO-d6	2.15(m) 3.2 to 3.4(m)	6.10	0.21 (d8)	4.14 (multi-plets)	4.74 4.75 (d5)	6.38 (s)	5.16 5.38 5.16 5.41 (J12)	3.38	$ \begin{array}{c} \text{CH}_3^m \\ \diagup \\ \text{O} \\ \parallel \\ \text{CHOCCH}_2\text{CH} \\ \quad \\ \text{CH}_3^j \quad \text{CH}_3^m \end{array} $	i 3.05(m) j 8.50(d6) k 7.75(d6) l 7.94(m) m 9.07(d7)
5	DMSO-d6	2.17(m) 3.2 to 3.4(m)	6.10	0.22 (d8)	4.12 4.15 (dd 8,5)	4.84 4.86 (d5)	6.26 6.49 (J18)	5.17 & 5.37 5.17 & 5.41 (J13)	3.40	$ \begin{array}{c} \text{O} \\ \parallel \\ \text{CHOCCH}_2 \\ \quad \\ \text{CH}_2\text{CH}_2\text{CH}_3 \\ k \quad l \quad m \end{array} $	i 3.05 & 3.15(16) j 7.94 k 8.20(m) l 8.3 to 8.8(m) m 9.09(17)
6	DMSO-d6	2.16(m) 3.2 to 3.4(m)	6.10	0.22 (d8)	4.12 4.14 (dd 8,5)	4.74 4.76 (d5)	6.28 6.49 (J18)	5.16 & 5.38 5.16 & 5.42 (J13)	3.38	$ \begin{array}{c} \text{O} \\ \parallel \\ \text{CHOCCH}_2 \\ \quad \\ \text{CH}_2\text{CH}_3 \\ k \quad l \end{array} $	i 3.12, 3.21(16) j 7.93 k 8.17(m) l 9.07(17)
7	DMSO-d6	2.14(s) 3.2 to 3.4(m)	6.06	0.20 (d8)	4.10 (m)	4.72 (d4)	6.36 (s)	5.15 5.36 (J12)	observed by a_2 and a_3	$ \begin{array}{c} \text{O} \\ \parallel \\ \text{CHOCCH}_2 \\ \quad \\ \text{CH}_2(\text{CH}_2)_4\text{CH}_3 \\ i \quad j \quad k \quad l \end{array} $	i 3.08(m) j 8.20(m) k 8.4 to 8.9(m) l 9.11(broad s) m 7.91(s)

TABLE 1 (Continued)

Example No.	Solvent	τ (100MHz; J Hz)								R	Assignments for R
		a_1, a_7, a_3	b	c	d	e	f	g	h		
9 Isomer A	DMSO-d ₆	2.16(d) 3.37 (dd 3,2) 3.30 (d 3)	6.10	0.21 (d8)	4.12 (dd 8,5)	4.74 (d5)	6.40 ABq	5.16 5.39 (J12)	3.40	$ \begin{array}{c} \text{O} \\ \parallel \\ \text{i} - \text{CHOCCH}_3 \\ \\ \text{CH}_3 \\ \text{j} \end{array} $	i 3.08(q5) j 8.51(d5) k 7.94
9 Isomer B	DMSO-d ₆	2.17(d2) 3.38 (dd 3,2) 3.29 (d 3)	6.09	0.22 (d8)	4.15 (dd 8,5)	4.76 (d5)	6.39 ABq	5.15 5.40 (J12)	3.40	$ \begin{array}{c} \text{O} \\ \parallel \\ \text{i} - \text{CHOC-CH}_3 \\ \\ \text{CH}_3 \\ \text{j} \end{array} $	i 2.99(q5) j 8.50(d5) k 7.94

TABLE 1 (Continued)

Example No.	Solvent	ν_{\max} (cm ⁻¹)				
		NH ₂ + NH	β -lactam	CONH	CO ₂ R	OCONH ₂
1	CHBr ₃	3540 3400	1780	1686 1520	1750 1730	1730 1584
2	CHBr ₃	3514 3380	1786	1686 1520	1750 1730	1730 1582
3	Nujol (Registered Trade Mark)	3470 3370 3300	1790	1680 1534	1750 1730	1720 1592
4	CHBr ₃	3500 3380	1790	1688 1522	1734 (strong)	1730 1588
5	CHBr ₃	3510 3380	1790	1688 1522	1758 1734	1730 1588
6	CHBr ₃	3520 3390	1788	1690 1520	1756 1732	1730 1584

TABLE I (Continued)

Example No.	Solvent	ν_{\max} (cm ⁻¹)				
		NH ₂ + NH	β -lactam	CONH	CO ₂ R	OCONH ₂
7	CHBr ₃	3502 3370	1782	1690 1518	1750 1730	1730
Isomer A 9	Nujol	3504 3440 to 3100	1780	1661 1521	1750	1710
Isomer B 9	Nujol	3420 3310 3290 3220	1778	1661 1528	1712	1710

Example A.

Tablet.

Composition:—

5	1-Acetoxyethyl (6R,7R) 3-carbamoyloxymethyl - 7 - [(Z)-2-(fur-2-yl)-2-methoxyiminoacetamido]-ceph-3-em-4-carboxylate (micronised)	326.0 mg
	Sodium starch glycolate (Primojel)	8.0 mg
10	Microcrystalline cellulose (Avicel* PH101)	64.0 mg
	Magnesium stearate	2.0 mg
	Total weight	400.0 mg

* The word "Avicel" is a registered Trade Mark.

Method of Preparation.

The magnesium stearate was blended with the active ingredient and tablet slugs were prepared by direct compression. The slugs were broken down through 12 mesh, 16 mesh and 20 mesh consecutively and the granules were blended with the sodium starch glycolate and microcrystalline cellulose. The blend was compressed on concave punches to a tablet weight of 400 mg. The tablets may be film coated by the aqueous or organic solvent method using cellulose derivatives with plasticisers and colouring matter. As an alternative to the preliminary slugging stage, the active ingredient may be densified by roller compaction.

Example B.

Powder for oral suspension (in sachet).

Composition (per sachet).

35	1-Acetoxyethyl (6R,7R)-3-carbamoyloxymethyl-7-[(Z)-(2-fur-2-yl)-2-methoxyiminoacetamido]-ceph-3-em-4-carboxylate (milled)	326.0 mg
40	Lecithin	25 mg
	Sodium carboxymethyl cellulose (low viscosity)	90 mg
	Spray-dried orange flavour	150 mg
	Caster sugar	2.2 g


Method of Preparation.

The lecithin was dissolved in chloroform and triturated with the active ingredient (previously milled using a fluid energy mill). The chloroform was allowed to evaporate and the resultant solid powdered. It was then blended intimately with the sodium carboxy-

methyl cellulose and the flavour. This blend was then further blended with the caster sugar adding the latter in two stages. It was intended that the correct weight should be filled into a sachet of suitable laminated foil and sealed by heat. The powder would be used by constituting with about 15 mls. water shortly before administration.

WHAT WE CLAIM IS:—

1. Compounds of the formula



(wherein R¹ is a primary or secondary alkyl group containing 1 to 4 carbon atoms and R² is a primary or secondary alkyl group containing 1 to 6 carbon atoms provided that at least one of the groups R¹ and R² is methyl).

2. A compound as claimed in claim 1 wherein R¹ is a methyl group and R² is an alkyl group containing 2 to 4 carbon atoms.

3. A compound as claimed in claim 1 wherein R² is a methyl group and R¹ is a primary or secondary alkyl group containing 1 to 4 carbon atoms.

4. 1 - Acetoxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

5. 1 - Propionyloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

6. 1 - Butyryloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

7. 1 - Isovaleryloxyethyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

8. 1 - Acetoxyheptyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.


9. 1 - Acetoxybutyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.

10. 1 - Acetoxypropyl (6R,7R) - 3 - carbamoyloxymethyl - 7 - [(Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetamido]ceph - 3 - em - 4 - carboxylate.


11. Diastereoisomers of compounds as claimed in any of the preceding claims.

12. A process for the preparation of a compound of formula I (as defined in claim 1) which comprises either (A) reacting (6R,

- 7R) - 3 - carbamoyloxymethyl - 7 - {(Z)-
2 - (fur - 2 - yl) - 2 - methoxyiminoacet-
amido]ceph - 3 - em - 4 - carboxylic acid
(i.e. cefuroxime) or a salt thereof with a
haloester of formula



(wherein R^1 and R^2 are as defined in claim 1 and X is halogen); or (B) acylating a compound of formula



- (wherein R^1 and R^2 are as defined in claim 1), or an acid addition salt of N-silyl derivative thereof, with (Z) - 2 - (fur - 2 - yl) - 2 - methoxyiminoacetate acid or a reaction derivative thereof.

13. A process as claimed in claim 12

wherein cefuroxime or a salt thereof is reacted with a haloester of formula (II) in which X is chlorine, bromine or iodine.

14. A process as claimed in claim 12 or claim 13 wherein an alkali metal or onium salt of cefuroxime is reacted with the said haloester of formula (II).

15. A process as claimed in claim 12 or claim 13 wherein cefuroxime is reacted with the said haloester of formula (II) in the presence of a base.

16. A process as claimed in claim 12 substantially as herein described.

17. A process for the preparation of compounds of formula I (as defined in claim 1) substantially as herein described in any of Examples 1 to 9.

18. Compounds of formula I (as defined in claim 1) whenever prepared by a process as claimed in any of claims 12 to 17.

19. A pharmacological composition comprising a compound as claimed in any of claims 1 to 11 in association with at least one pharmaceutical carrier or excipient.

FRANK B. DEHN & CO.,
Chartered Patent Agents,
Imperial House, 15-19, Kingsway,
London, W.C.2.